

Momentum-Resolved Inelastic X-ray Scattering as a Novel Tool to Study Charge Gap in Complex Insulators

M. Z. Hasan¹, E. D. Isaacs², Z.-X. Shen¹, L. L. Miller³

¹*Department of Applied Physics and Stanford Synchrotron Radiation Laboratory (SSRL) of Stanford Linear Accelerator Center (SLAC), Stanford University, Stanford, CA 94305*

²*Bell Laboratories, Lucent Technologies, Murray Hill, NJ 07974*

³*Department of Physics, Iowa State University and Ames Laboratory, Ames, IA 50011*

(February 1, 2008)

We report particle-hole pair excitations in a cuprate insulator in the intermediate regimes of momentum-transfers using high energy inelastic x-ray scattering. The excitation spectra show dispersive features near the Mott edge which shed light on the momentum structure of the upper Hubbard band in cuprates. We briefly discuss the potential use of such a technique to study the momentum dependence of unoccupied bands and $\mathbf{q}$ -dependent charge fluctuations in complex insulators.

After several decades of research efforts, electronic structure of late transition metal oxides lacks comprehensive understanding. The existence of exotic electronic, magnetic and optical properties such as high T_c superconductivity as exhibited by the cuprates or colossal magnetoresistance as in the manganites or highly nonlinear optical responses as observed in the nickelates are believed to be related to the strong electron-electron Coulomb correlations in these systems [1-4]. This suggests the importance of a thorough study of their correlated charge dynamics. Angle-resolved photoemission (ARPES) which probes the occupied electronic states in a momentum-resolved manner has been successful in characterizing the electronic structure of these oxides [4-6] whereas electron-energy-loss spectroscopy (EELS) is limited to low momentum transfers due to multiple scattering effects arising from its strong coupling nature [7] and being surface sensitive both ARPES and EELS require extensive ultra-high vacuum (UHV) sample preparation [4,7]. Any bulk study of the momentum-resolved electronic structure specially the unoccupied bands is absent for these correlated insulators.

In last several years, with the advent of high brightness synchrotron facilities, inelastic x-ray scattering has been developing as a tool to study the BULK electronic structure of condensed matter systems [4-10]. X-ray scattering from the valence charge distribution is fairly weak thus difficult to distinguish from the total scattering signal especially in high-Z materials making such experiments quite difficult to perform. Recent experimental and theoretical investigations have shown that by tuning the incident energy near an x-ray absorption edge a large enhancement can be achieved making the study of valence excitations feasible in high-Z systems [5,7-10].

The observation of a low-energy charge-transfer gap has been reported recently with nonzero- $\mathbf{q}$ in a parent cuprate using inelastic x-ray scattering [13,16] and has extensively been studied by optical spectroscopies (with $q \sim 0$ momentum transfer) [17]. A similar excitation band is also seen in EELS at low momentum transfers (< 1

$\text{\AA}^{-1}$) [7]. In this paper, we report some novel applications of inelastic x-ray scattering and discuss the information it can provide about electronic excitations in solids.

Experiments were performed at the X-21-A3 wiggler beamline at the National Synchrotron Light Source of Brookhaven National Laboratory with an overall energy resolution of 0.4 eV determined by fitting the elastic scattering and typical inelastic count rates from the sample were 20 to 30 counts per minute at energy losses of several electron volts around 250 mA ring current. The scattered light was reflected from a germanium analyzer and focussed onto a solid-state detector. Energy analysis was done by rotating the analyzer and translating the detector accordingly at the focus of the analyzer. Incident energy was kept fixed near the Cu K-edge ($E_0 = 8995.8$ eV) determined from the fluorescence profile. The experiment was performed in horizontal scattering geometry where $\mathbf{q}$ was varied in the plane defined by the incident polarization. Background was measured by keeping track of scattering intensities on the energy gain side (several eV on the energy gain spectrum) which was about 1-2 counts per minute. The $\text{Ca}_2\text{CuO}_2\text{Cl}_2$ crystals used for this experiment were grown and characterized by techniques described previously [18]. Due to its extreme hygroscopic nature the crystal used for the experiment was always kept, prepared and preoriented for the mount under dry N_2 and chemical desiccant environments and the experiment was performed in a vacuum system with transparent capton windows.

Fig. 1(A) shows inelastic x-ray scattering spectra near Cu K-edge from parent cuprate $\text{Ca}_2\text{CuO}_2\text{Cl}_2$ with varying momentum transfers along the $\langle 110 \rangle$ direction (45 degrees to the Cu-O bond direction) and Fig. 1(B) shows spectra with momentum transfers along the $\langle 100 \rangle$ direction (the Cu-O bond direction). Elastic scattering has been removed by fitting and all the spectra in each panel were normalized near 8 eV energy-loss so the intensities reported here are relative. Each spectrum exhibits two features - one broad excitation band around 5.8 eV and a weak feature around 2.45 to 3.9 eV for different $\mathbf{q}$ -values.




FIG. 1. Inelastic x-ray scattering spectra near Cu K-edge are shown along two directions in the Cu-O plane : (A) Scattering with $\mathbf{q}$ along the $\langle 110 \rangle$ -direction (45 degrees to the Cu-O bond direction). The values of $\mathbf{q}$ for the spectra bottom to top are $(1.9\pi, 1.9\pi)$, $(1.7\pi, 1.7\pi)$, $(1.5\pi, 1.5\pi)$, $(1.2\pi, 1.2\pi)$, $(1.1\pi, 1.1\pi)$ respectively. (B) Scattering with $\mathbf{q}$ along the $\langle 100 \rangle$ -direction (the Cu-O bond direction). The values of $\mathbf{q}$ for the spectra bottom to top are $(2.1\pi, 0)$, $(2.2\pi, 0)$, $(2.5\pi, 0)$, $(2.7\pi, 0)$, $(3.1\pi, 0)$ respectively. Incident photon energy $E_o = 8995.8$ eV.

The broad feature centered around 5.8 eV changes shape with change of $\mathbf{q}$ (also the associated changes in polarization) but does not show any significant dispersion (compared to its width) in either of the directions. Nondispersive behavior exhibited by the feature for a change of $\mathbf{q}$ over a large range (1.4π to 3.1π) strongly suggests that it has a highly local character. Based on electronic structure calculations the 5.8-eV feature is believed to be a charge transfer excitation from an occupied state with b_{1g} symmetry to an empty state with a_{1g} symmetry [13,19]. The lower energy feature, on the other hand, has a significant movement in changing $\mathbf{q}$ from $(1.9\pi, 1.9\pi)$ to $(1.1\pi, 1.1\pi)$ as seen in Fig. 1(A). The feature disperses upward about 1.4 eV monotonically in this direction. Whereas if the momentum transfer is along the bond direction it does not show much dispersion in going from $(2.1\pi, 0)$ to $(2.5\pi, 0)$. But in going from $(2.5\pi, 0)$ to $(3.1\pi, 0)$ it disperses upward by about 0.5 eV. Dispersion behavior in two directions are compared in fig-2(A) where we plotted the center of gravity of the spectral weights of the low-energy feature. Overall dispersion along the bond direction is much weaker than dispersion along the diagonal direction in the Cu-O lattice.

Inelastic x-ray scattering measures the dynamical charge-charge correlation function (charge fluctuations) which can be interpreted as particle-hole pair excitations in the range of momentum-transfers comparable to the size of the equivalent Brillouin zones of the system [13-15]. Near an absorption edge the measured dynamical

response function gets modified due to the presence of a core hole and an excited electron (hence sensitivity to polarization of the incident x-ray) but it can still be interpreted as composites of pair excitations [13-15]. In the scattering process, the core-hole created by the X-ray photon near the absorption edge causes electronic excitations in the valence band which creates a hole in the occupied band and promotes an electron to the unoccupied band across the charge gap in an insulator. A scenario for the origin of the lower energy feature observed in our experimental data can consist of charge excitations across the characteristic gap in the system - the charge-transfer gap (effective Mott gap) which is related to the gap seen in optical experiments (~ 2 eV) [17].

In an effective single-particle band picture, one such possibility [15] is shown in Fig-2(B), it is an excitation from the occupied band to the unoccupied band. Particle-hole pairs created through the scattering process absorb the energy and momentum lost from the X-ray photons [14,15]. Scanning the feature in $(\omega, \mathbf{q})$ space reveals the detailed dynamics of the particle-hole pair in the system. From a series of ARPES experiments we know that the occupied band has a maximum near $\mathbf{k} = (\pi/2, \pi/2)$ [4-6]. In a single-electron-like band picture, if we assume that the unoccupied band as a mirror image




FIG. 2. The momentum dependence of the center of gravity of the low-energy inelastic feature is compared along the $\langle 110 \rangle$ direction and the $\langle 100 \rangle$ direction and shown in (A). Relative excitation energies are plotted referenced to the energy (E_o) at $\mathbf{q} = (0,0)$ along each direction. (B) shows one possible schematic cartoon of energy-gap in the electronic structure of the insulators which separates the occupied states from the unoccupied ones. There exists a charge excitation gap between the occupied band and the unoccupied upper Hubbard band in cuprate insulators. The unique advantage of high energy inelastic x-ray scattering is that it can probe charge excitations with finite momentum transfers over a large portion of the Brillouin Zones. In insulating cuprates these excitations reveal the momentum structure of the upper Hubbard band.

of the occupied band across the gap, in a simplistic view, we may expect a folding in dispersion of particle-hole pair excitations near $\mathbf{q} \sim (1.5\pi, 1.5\pi)$. Instead, we observe in the data a clear monotonic increase of the excitation gap along $(1.9\pi, 1.9\pi)$ to $(1.1\pi, 1.1\pi)$ and flatness from $(2.1\pi, 0)$ to $(2.5\pi, 0)$ and then monotonic increase again in going from $(2.5\pi, 0)$ to $(3.1\pi, 0)$. This dispersive behavior is an indication that charge-transfer or effective Mott excitations can not be viewed as composite single-particle excitations but should rather be viewed as a single correlated two-body excitation as argued by Tsutsui et.al. [15].

We also note that in the case of a hole having a Zhang-Rice singlet wavefunction [20] forming a pair with an electron (at the upper Hubbard band) would have different propagation probabilities along the Cu-O bond and 45 degrees to the Cu-O bond due to the antiferromagnetic correlation of the lattice. This difference can be relatively significant because of the large exchange interactions (a consequence of large Hubbard-U) exhibited by the undoped cuprate insulator. Dispersions observed in the experiment reflect the nature of the dispersive Hubbard bands and we can estimate the relevant bandwidth involved in the excitation process. The single-hole bandwidth in this system is fairly narrow (~ 0.35 eV) as known from ARPES [4-6]. We speculate that the coherent dispersion of the particle-hole pair (~ 1.4 eV) gains most contribution from the electron bandwidth on the upper Hubbard band. These results are consistent with numerical studies of Hubbard hamiltonian with long-range hopping [15]. Flatness along the bond direction, as seen in the data, is probably due to the fact that the lowest-energy state of the upper Hubbard band is shifted from $\mathbf{k} = (\pi/2, \pi/2)$ [15,16]. We postpone the extraction of quantitative details for the higher resolution experiments in near future.

These results demonstrate the feasibility of inelastic x-ray scattering to study electronic excitations at the gap edge of the Mott insulators and its potential in gaining important information about correlated electron systems. Momentum-resolved nature of the upper Hubbard band of charge-transfer or Mott insulators (parent high T_c and CMR compounds) has so far been unknown but such information is crucial in understanding the fascinating Mott phenomena in a variety of other condensed matter systems. The anisotropy and direct/indirect nature of Mott gap can be studied in other insulating systems such as manganites in connection to CMR physics or Kondo insulators in connection to their non-Fermi liquid behavior. There have been many speculations about the possibility of highly $\mathbf{q}$ -dependent charge fluctuations such as the dynamical charge stripes in high T_c cuprates [21,22]. Very high resolution inelastic x-ray scattering could possibly be used to study dynamic stripes because of the direct coupling of x-rays to the electronic charges (unlike neutrons which has so far been used to probe dynamical correlations either of the lattice or the spin directly). Similarly, the fluctuations of the orbitally or-

dered phases (orbital waves/orbitons) in orbitally degenerate systems such as manganese oxides can be studied using polarization dependent inelastic x-ray scattering near an atomic-core resonance[23]. Particle-hole excitations, in general, are fundamental to the transport phenomena so it is of importance to use two-particle spectroscopies in a momentum-resolved mode so that an understanding of the nonlocal and anisotropic interaction potentials can be developed which determine various groundstates of a correlated system. With the availability of high brightness synchrotron facilities a new frontier of understanding correlated systems would emerge through utilizing this class of scattering techniques.

We gratefully acknowledge P. Abbamonte, K. Tsutsui and C. Kao for useful suggestions. This work performed at the National Synchrotron Light Source of Brookhaven National Laboratory was jointly supported by the Department of Energy (BES/MSD) through Stanford Synchrotron Radiation Lab of Stanford Linear Accelerator Center, Stanford, California and Bell-Laboratories of Lucent Technologies, New Jersey.

-
- [1] N. F. Mott Proc.Phys.Soc.London A **62**, 416 (1949).
 - [2] J. Hubbard Proc.Phys.Soc.London A **277**, 237 (1964).
 - [3] J. Zaanen *et. al.*, Phys. Rev. Lett. **55**, 418 (1985).
 - [4] Z. X. Shen *et. al.* Science **267**, 343 (1995).
 - [5] B. O. Wells *et. al.*, Phys. Rev. Lett. **74**, 964 (1995).
 - [6] F. Ronning *et. al.*, Science **282**, 2067, (1998).
 - [7] Y. Y. Wang *et. al.*, Phys. Rev. Lett. **77**, 1809 (1997).
 - [8] C. C. Kao *et. al.*, Phys. Rev. B **54**, 16 361 (1996)
 - [9] F. Sette *et. al.*, Phys. Rev. Lett. **77**, 83 (1996).
 - [10] E. D. Isaacs *et. al.*, Rev. Lett. **76**, 4211 (1996).
 - [11] M. H. Krisch *et. al.*, Phys. Rev. Lett. **78**, 2843 (1997).
 - [12] J. P. Hill *et. al.*, Phys. Rev. Lett. **80**, 4967 (1998).
 - [13] P. Abbamonte *et. al.*, Phys. Rev. Lett. **83**, 860 (1999).
 - [14] P. M. Platzman *et. al.*, Phys. Rev. B **57**, 11 107 (1998).
 - [15] K. Tsutsui *et. al.*, Phys. Rev. Lett. **83**, 3705 (1999).
 - [16] M. Z. Hasan *et. al.*, Science **288**, 1811 (2000)
 - [17] S. L. Cooper *et. al.*, Phys. Rev. B **47**, 8233 (1993)
 - [18] L. L. Miller *et. al.*, Phys. Rev. B **41**, 1921 (1990).
 - [19] M. E. Simon *et. al.*, Phys. Rev. B **54**, R3780 (1996).
 - [20] F. C. Zhang *et. al.*, Phys. Rev. B **37**, 3759 (1988).
 - [21] J. M. Tranquada *et. al.*, Nature **375**, 561 (1995).
 - [22] S. A. Kivelson *et. al.*, Nature **393**, 550 (1998).
 - [23] H. Kondo *et. al.*, cond-mat/0012353 (2000)